

Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at http://about.jstor.org/participate-jstor/individuals/early-journal-content.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.



G. Edwards delineavit A: 1770.

Baire ..

V. Description of a Bird from the East Indies; in a Letter to James West, Esq; President of the Royal Society; from Mr. George Edwards, F. R. S.

SIR,

Read Jan. 17, N August last, a friend of mine carried me with him to Valentine House, near Ilford in Essex, the seat of Charles Raymond, Esq; to see some curious birds and other animals, from the East Indies; amongst these, I discovered a rare bird, not before known to me *. It is of a new genus, and the only species of the genus hitherto come to my knowledge. It is about the bigness of a heron [see Tab. II.]; and has a good deal of the appearance of birds of the heron and crane kind, except that the neck is a little shorter. On first sight, I thought the bird belonged to that genus; but, on a closer view, I judged it to be no wader in the water, for though the legs be as long or longer than in herons, &c. yet they are feathered down to the knees, which we do not find in birds who wade in shallow waters, to seek their

^{*} This bird was described, under the name of the Sagittarius from the Cape of good Hope, by Mr. Vosmaer, keeper of the Statholder's Museum at the Hague, in one of his publications in low Dutch, printed at Amsterdam, 1769, in 4to, with a coloured cut of the said bird. It seems to feed equally on sless and sish; which accounts for his uniting the characters of birds of prey, and of waders in water. M. M.

food. The toes in this bird are also much shorter than they are in herons, so that I think it must be placed amongst land birds. The bill is exactly like those of hawks, and other birds of prey, which is the only instance I have discovered in any of the long legged kind of birds; the talons or claws are small and unfit for a bird of prey, and the eyes are of a dark colour placed in spaces covered with a bare skin of an orange colour, on each fide of the head. It hath a beautiful crest composed of many long painted feathers tipped with black hanging backward. The beak, head, neck, back, breaft, and upper covert feathers of the wings are of a blueish ash colour, rather lighter on the breast than on the The belly, thighs, the greater wing-feathers, and tail, are black, the tail feathers being tipped with white; the legs and feet are of a reddish flesh colour, the claws black. This bird was called a fnake-eater, by those who brought it from India. believe it may prey on small serpents, lizards, and other small reptiles. Another bird was brought with this, supposed to be the male of this species, which died foon after it was landed Mr. Raymond's fervant told me that it was fomething larger, and the crest longer, the head black, but that in other respects the two birds agreed.

I am, SIR,

Your most humble servant,

College of Physicians, January, 1771.

Geo. Edwards.